For Research Use Only

MCM4 Monoclonal antibody, PBS Only



Catalog Number: 67103-1-PBS

Basic Information

Catalog Number: 67103-1-PBS

Size: 1 mg/ml Source: Mouse Isotype:

lgG2b Immunogen Catalog Number:

AG28642

Tested Applications:

WB,Indirect ELISA Species Specificity: Human, mouse, rat

GenBank Accession Number:

BC031061 GeneID (NCBI): 4173

UNIPROT ID: P33991 Full Name:

minichromosome maintenance

complex component 4

Calculated MW: 863 aa, 97 kDa Observed MW: 97 kDa

Purification Method: Protein A purification

CloneNo.: 2H2A1

Applications

Background Information

DNA replication licensing factor MCM4 (MCM4), also named Cdc21, acts as component of the MCM2-7 complex (MCM $complex) which is the putative replicative helicase essential for 'once per cell cycle' \, DNA \, replication initiation \, and \, complex \, and \, complex \, are the putative replicative helicase essential for 'once per cell cycle' \, DNA \, replication initiation \, and \, complex \, are the putative replicative helicase essential for 'once per cell cycle' \, DNA \, replication initiation \, and \, complex \, are the putative replicative helicase essential for 'once per cell cycle' \, DNA \, replication initiation \, and \, complex \, are the putative replicative helicase essential for 'once per cell cycle' \, DNA \, replication initiation \, and \, complex \, are the putative replication in the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, and \, complex \, are the putative replication \, are the putative replication \, and \, complex \, are the putative replication \, are the putative replication \, are the p$ elongation in eukaryotic cells. The active ATPase sites in the MCM2-7 ring are formed through the interaction surfaces of two neighboring subunits such that a critical structure of a conserved arginine finger motif is provided in trans relative to the ATP-binding site of the Walker A box of the adjacent subunit. The six ATPase active sites, however, are likely to contribute differentially to the complex helicase activity.

Storage


Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer: PBS Only

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67103-1-lg (MCM4 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 67103-1-PBS in a different storage buffer formulation.