For Research Use Only

HOXA7 Monoclonal antibody, PBS Only



Catalog Number: 67112-1-PBS

Basic Information

Catalog Number:

67112-1-PBS

Size: 1mg/ml

Source: Mouse Isotype: IgG1

Immunogen Catalog Number:

AG19025

025 230 aa, 25 kDa

Observed MW:

20 kDa

BC148692

3204

P31268 Full Name:

GeneID (NCBI):

UNIPROT ID:

homeobox A7

Calculated MW:

GenBank Accession Number:

Purification Method:

Protein A purification

CloneNo.: 2A7H6

Applications

Tested Applications: WB, IF, ELISA Species Specificity:

Human

Background Information

HOX genes play a fundamental role in the development of the vertebrate central nervous system, heart, axial skeleton, limbs, gut, urogenital tract and external genitalia. The homeobox gene Hoxa-1 is transcriptionally regulated by retinoic acid (RA) and encodes a transcription factor, which has been shown to play important roles in cell differentiation and embryogenesis. Hoxa-1 is also expressed in cancers, such as mammary tumors, though it is not expressed in normal gland or in precancerous mammary tissues. At embryonic stages, Hoxa-2 is expressed in the mesenchyme and epithelial cells of palate, however its expression is restricted to the tips of the growing palatal shelves. Hoxa-2 protein is predominantly expressed in the nuclei of cells in the ventral mantle region of the developing embryo. In the developing and adult mouse spinal cord, Hoxa-2 protein may contribute to dorsal-ventral patterning and/or to the specification of neuronal phenotype. Hoxa-7 functions as a potent transcriptional repressor and its action as such requires several domains, including both activator and repressor regions. Hoxa-7 is expressed in the fetal liver, lung, skeletal muscle, kidney, pancreas and placenta.

Storage


Storage:

Store at -80°C.


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C Storage Buffer.

PBS only

Selected Validation Data



THP-1 cells were subjected to SDS PAGE followed by western blot with 67112-1-1g (HOXA7 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 67112-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using 67112-1-1g (HOXA7 antibody) at dilution of 1:100 and CoraLite488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 67112-1-PBS in a different storage buffer formulation.