For Research Use Only

## FLNA Monoclonal antibody, PBS Only

Catalog Number: 67133-1-PBS



**Basic Information** 

Catalog Number:

GenBank Accession Number:

Purification Method:

67133-1-PBS

BC014654
GeneID (NCBI):

Protein A purification

Size: 1 mg/ml

GenelD (NCBI) 2316 CloneNo.: 1G4H3

Source: Mouse

UNIPROT ID: P21333 Full Name:

Isotype: IgG2a

filamin A, alpha (actin binding

protein 280)

AG23291

Immunogen Catalog Number:

Calculated MW: 2647 aa, 280 kDa

Observed MW:

280 kDa

**Applications** 

Tested Applications:

WB,IHC,IF,ELISA

Species Specificity:

**Background Information** 

FLNA encodes a 280 kDa widely expressed actin binding protein called filamin A which can crosslinks actin filaments and links actin filaments to membrane glycoproteins to form a three-dimensional network (PMID:29931263). And filamin A could interact with many other proteins, involved in receptor activation, cell migration and adhesion, cell proliferation, inflammation and tumorigenesis, studies have been reported that FLNA overexpressed in multiple types of tumors, suggesting FLNA may involved in cancer aggressiveness(PMID:29100390). What's more, FLNA is an causative gene of periventricular nodular heterotopia (PNH) (PMID: 29062687)

Storage

Storage:


Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C  $\,$ 

Storage Buffer:

PBS only


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67133-1-lg (FLNA antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 67133-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human cervical cancer tissue slide using 67133-1-Ig (FLNA antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67133-1-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (-20°C Methanol) fixed Hela cells using FLNA antibody (67133-1-lg, Clone: 1G4H3) at dilution of 1:800 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 67133-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Ethanol) fixed Hela cells using FLNA antibody (67133-1-lg, Clone: 1G4H3) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 67133-1-PBS in a different storage buffer formulation.



MCF-7 cells were subjected to SDS PAGE followed by western blot with 67133-1-lg (FLNA antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 67133-1-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 67133-1-Ig (FLNA antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 67133-1-PBS in a different storage buffer formulation.