For Research Use Only

SUMO2/3 Monoclonal antibody, PBS Only



Purification Method:

Protein A purification

CloneNo.:

1A1B3

Catalog Number: 67154-1-PBS

Featured Product

Basic Information

Catalog Number: 67154-1-PBS

Size:
1mg/ml
Source:
Mouse

Isotype:Full Name:IgG2bSMT3 suppressor of mif two 3

Immunogen Catalog Number:

AG28672

homolog 2 (S. cerevisiae)

Calculated MW:

11 kDa

BC008450

6613

P61956

GeneID (NCBI):

UNIPROT ID:

GenBank Accession Number:

Observed MW: 18 kDa

Applications

Tested Applications:

WB, IHC, IF/ICC, Indirect ELISA

Species Specificity: human, mouse, rat

Background Information

Ubiquitin is most famous for its function in targeting proteins for degradation by the 26S proteasome, ubiquitin needs to be attached to a substrate in chains (polyubiquitylation) before being recognized by proteasome. Similarly, SUMO (small ubiquitin-related modifier) can be linked to substrates in chains (polysumoylation), SUMO modification has been implicated in many important cellular processes including the control of genome stability, signal transduction, targeting to and formation of nuclear compartments, cell cycle and meiosis. There are 4 confirmed SUMO isoforms in human, SUMO-1, SUMO-2, SUMO-3 and SUMO-4. SUMO-3 are nearly identical but are distinct from SUMO-1. SUMO2/3 conjugation was recently widely involved in neuroprotective activities. A substitution (M55V) of SUMO4 was strongly associated with the pathogenesis of type 1 diabetes (T1D) involving NF kappa B related mechanisms.

Storage


Storage:

Store at -80°C.


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C Storage Buffer:

PBS Only


Selected Validation Data




WB result of SUMO2/3 antibody (67154-1-lg; 1:3000; incubated at room temperature for 1.5 hours) with sh-Control and sh-SUMO2/3 transfected HEK-293 cells. This data was developed using the same antibody clone with 67154-1-PBS in a different storage buffer formulation.




Various lysates were subjected to SDS PAGE followed by western blot with 67154-1-1g (SUMO 2/3 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 67154-1-PBS in a different storage buffer formulation




Immunohistochemical analysis of paraffinembedded human prostate cancer tissue slide using 67154-1-Ig (SUMO2/3 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67154-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human prostate cancer tissue slide using 67154-1-1g (SUMO2/3 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67154-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using SUMO2/3 antibody (67154-1-lg, Clone: 1A1B3) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1), CL594-phalloidin (red). This data was developed using the same antibody clone with 67154-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using SUMO2/3 antibody (67154-1-lg, Clone: 1A1B3) at dilution of 1:800 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 67154-1-PBS in a different storage buffer formulation.