## For Research Use Only

## HIP1 Monoclonal antibody, PBS Only

Catalog Number: 67163-1-PBS **Featured Product** 



**Basic Information** 

Catalog Number:

GenBank Accession Number: BC110545

**Purification Method:** Protein G purification

67163-1-PBS

GeneID (NCBI):

Size: 1 mg/ml

Isotype: lgG1

3092

CloneNo.: 3D3G3

Source: Mouse

**UNIPROT ID:** 000291 Full Name:

huntingtin interacting protein 1

Immunogen Catalog Number:

Calculated MW: 1037 aa, 116 kDa

AG17982

Observed MW:

110-120 kDa

**Applications** 

**Tested Applications:** 

WB,Indirect ELISA,IHC,IF

Species Specificity:

Human, mouse, pig

## **Background Information**

HIP1 (huntingtin interacting protein-1) is a endocytic protein that plays a role in clathrin-mediated endocytosis and trafficking. HIP1 can interact with huntingtin (HTT) protein whose mutations cause Huntington's disease (HD), thus gets involved in neurodegeneration. By stabilizing receptor tyrosine kinases following ligand-induced endocytosis, HIP1 may act to promote cell survival. HIP1 is overexpressed in a variety of human cancers and may be a potential predictor for cancer progression and prognosis.

Storage


Storage:

Store at -80°C.


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer: PBS Only


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67163-1-lg (HIP1 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 67163-1-PBS in a different storage buffer formulation.




WB result of HIP1 antibody (67163-1-lg; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-HIP1 transfected Hela cells. This data was developed using the same antibody clone with 67163-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 67163-1-lg (HIP1 antibody) at dilution of 1:300 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67163-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 67163-1-lg (HIP1 antibody) at dilution of 1:300 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67163-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed human colon cancer tissue using HIP1 antibody (67163-1-Ig, Clone: 3D3G3) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 67163-1-PBS in a different storage buffer formulation.