For Research Use Only

AMPK Gamma 1 Monoclonal antibody, PBS Only



Catalog Number:67182-1-PBS

Featured Product

Basic Information

Catalog Number: 67182-1-PBS

Size:
1 mg/ml
Source:
Mouse
Isotype:

Immunogen Catalog Number:

AG28907

lgG1

GenBank Accession Number: BC000358

GeneID (NCBI): 5571 UNIPROT ID:

P54619

Full Name: protein kinase, AMP-activated, gamma 1 non-catalytic subunit

Calculated MW: 38 kDa Observed MW: 36-38 kDa Purification Method:

Protein G purification CloneNo.: 1E11D4

Applications

Tested Applications: WB,Indirect ELISA Species Specificity: Human, mouse, rat

Background Information

Protein kinase, AMP-activated, gamma 1 non-catalytic subunit (PRKAG1, synonyms: AMPKG, MGC8666) is a regulatory subunit of the AMP-activated protein kinase (AMPK). AMPK is a heterotrimer consisting of an catalytic subunit, and non-catalytic and subunits. AMPK is an important energy-sensing enzyme that monitors cellular energy status. In response to cellular metabolic stresses, AMPK is activated, and thus phosphorylates and iAMP-activated protein kinase (AMPK) is a highly conserved heterotrimeric serine/threonine kinase widely characterised as a sensor of cellular energetic stress. AMPK is a heterotrimeric complex consisting of a catalytic α -subunit and two regulatory subunits (β and γ). AMPK is an important energy-sensing enzyme that monitors cellular energy status. In response to cellular metabolic stresses, AMPK is activated, and thus phosphorylates and inactivates acetyl-CoA carboxylase (ACC) and beta-hydroxy beta-methylglutaryl-CoA reductase (HMGCR), key enzymes involved in regulating de novo biosynthesis of fatty acid and cholesterol. AMPK gamma 1 is one of the gamma regulatory subunits of AMPK.

Storage


Storage:


Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C Storage Buffer.

PBS Only

Selected Validation Data





Various lysates were subjected to SDS PAGE followed by western blot with 67182-1-lg (AMPK gamma 1 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Alpha Tubulin Monoclonal antibody (HRP-66031) as loading control. This data was developed using the same antibody clone with 67182-1-PBS in a different storage buffer formulation.

WB result of AMPK Gamma 1 antibody (67182-1-1g; 1:20000; incubated at room temperature for 1.5 hours) with sh-Control and sh-AMPK Gamma 1 transfected HEK-293 cells. This data was developed using the same antibody clone with 67182-1-PBS in a different storage buffer formulation.