

# VDR Monoclonal antibody, PBS Only

Catalog Number: 67192-1-PBS

Featured Product

## Basic Information

**Catalog Number:**

67192-1-PBS

**Size:**

1 mg/ml

**Source:**

Mouse

**Isotype:**

IgG2a

**Immunogen Catalog Number:**

AG28188

**GenBank Accession Number:**

BC060832

**GeneID (NCBI):**

7421

**UNIPROT ID:**

P11473

**Full Name:**

vitamin D (1,25- dihydroxyvitamin D3) receptor

**Calculated MW:**

48 kDa

**Observed MW:**

48-55 kDa

**Purification Method:**

Protein A purification

**CloneNo.:**

1A9C1

## Applications

**Tested Applications:**

WB, Indirect ELISA

**Species Specificity:**

Human, mouse, rat

## Background Information

The vitamin D receptor (VDR), also known as NR111 (nuclear receptor subfamily 1, group I, member 1), is a member of the nuclear receptor family of transcription factors. Upon activation by vitamin D, the VDR forms a heterodimer with the retinoid-X receptor and binds to hormone response elements on DNA resulting in expression or trans-repression of specific gene products. It is an intracellular hormone receptor that specifically binds 1,25(OH)<sub>2</sub>D<sub>3</sub> and mediates its effects. Downstream targets of this nuclear hormone receptor are principally involved in mineral metabolism though the receptor regulates a variety of other metabolic pathways, such as those involved in the immune response and cancer. Defects in VDR are the cause of rickets vitamin D-dependent type 2A (VDDR2A). A disorder of vitamin D metabolism results in severe rickets, hypocalcemia and secondary hyperparathyroidism. Most patients have total alopecia in addition to rickets. The VDR exists two isoform with the MV 48 kDa and 54 kDa.

## Storage

**Storage:**

Store at -80°C.

**The product is shipped with ice packs. Upon receipt, store it immediately at -80°C****Storage Buffer:**

PBS Only


For technical support and original validation data for this product please contact:

T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)W: [ptgcn.com](http://ptgcn.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

# Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67192-1-Ig (VDR antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 67192-1-PBS in a different storage buffer formulation.