

For Research Use Only

CDK9 Monoclonal antibody

Catalog Number: 67256-1-Ig



Basic Information

Catalog Number: 67256-1-Ig	GenBank Accession Number: BC001968	Purification Method: Protein A purification
Size: 1000 µg/ml	GeneID (NCBI): 1025	CloneNo.: 1G6B3
Source: Mouse	UNIPROT ID: P50750	Recommended Dilutions: WB 1:5000-1:50000 IF 1:500-1:2000
Isotype: IgG1	Full Name: cyclin-dependent kinase 9	
Immunogen Catalog Number: AG29178	Calculated MW: 372 aa, 43 kDa	
	Observed MW: 38-42 kDa, 55 kDa	

Applications

Tested Applications: IF/ICC, WB, ELISA	Positive Controls:
Species Specificity: Human, Mouse, Rat	WB : HeLa cells, T-47D cells, HCT 116 cells, A431 cells, MOLT-4 cells, TF-1 cells, HSC-T6 cells, ROS1728 cells
	IF : HeLa cells,

Background Information

CDK9(Cyclin-dependent kinase 9) is a member of the Cdc2-like family of kinases. Its cyclin partners are members of the family of cyclin T (T1, T2a and T2b) and cyclin K. The CDK9/cyclin T complexes appear to be involved in regulating several physiological processes. CDK9 has also been described as the kinase of the TAK complex, which is homologous to the P-TEFb complex and involved in HIV replication. In addition, CDK9 seems to have an anti-apoptotic function in monocytes, that may be related to its control over differentiation of monocytes (PMID: 12432243). CDK9 has two isoforms with the molecular mass of 42 kDa and 55 kDa, and the relative abundance of Cdk9(42kDa) and Cdk9(55kDa) changes in different cell types (PMID: 12706900, 15780980).

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.1% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67256-1-Ig (CDK9 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using CDK9 antibody (67256-1-Ig, Clone: 1G6B3) at dilution of 1:1000 and Multi-rAbCoraLite Plus 594-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (RGAM004, 1:500) were used for detection.