

For Research Use Only

# ODC1 Monoclonal antibody

Catalog Number: 67336-1-Ig **Featured Product**



## Basic Information

Catalog Number:

67336-1-Ig

Size:

2000 µg/ml

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG10927

GenBank Accession Number:

BC025296

GeneID (NCBI):

4953

UNIPROT ID:

P11926

Full Name:

ornithine decarboxylase 1

Calculated MW:

461 aa, 51 kDa

Observed MW:

51 kDa

Purification Method:

Protein G purification

CloneNo.:

2A1E3

Recommended Dilutions:

WB 1:5000-1:50000

## Applications

Tested Applications:

WB, ELISA

Species Specificity:

Human

Positive Controls:

WB : MCF-7 cells, HUVEC cells, human placenta tissue

## Background Information

Ornithine decarboxylase (ODC) is also named as ODC1 and belongs to the Orn/Lys/Arg decarboxylase class-II family. It catalyzes the conversion of ornithine to putrescine, the first step and a major site of regulation of polyamine biosynthesis. The level of ODC is known to be controlled at several sites, namely transcription, translation, and enzyme degradation. Polyamines can stimulate the degradation of ODC as a type of negative feedback control (PMID:8486633). This protein can be phosphorylated in vivo (PMID:8798774). ODC1 can form a homodimer and only the dimer is catalytically active, as the active sites are constructed of residues from both monomers (PMID: 10623504). The molecular mass of ODC1 is 51 kDa, and the homodimer is 106 kDa.

## Storage

Storage:

Store at -20°C.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**

## Selected Validation Data



WB result of ODC1 antibody (67336-1-Ig; 1:30000; incubated at room temperature for 1.5 hours) with sh-Control and sh-ODC1 transfected MCF-7 cells.



MCF-7 cells were subjected to SDS PAGE followed by western blot with 67336-1-Ig (ODC1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.