## For Research Use Only

## RBM39 Monoclonal antibody

Catalog Number: 67420-1-Ig



**Basic Information** 

Catalog Number: GenBank Accession Number:

67420-1-lg BC141835
Size: GeneID (NCBI):
1000 μg/ml 9584
Source: UNIPROT ID:
Mouse Q14498
Isotype: Full Name:

IgG2b RNA binding motif protein 39

Immunogen Catalog Number:Calculated MW:AG15766530 aa, 59 kDaObserved MW:

60-66 kDa

Purification Method: Protein A purification

Protein A purifica CloneNo.: 2D2C8

Recommended Dilutions: WB 1:5000-1:50000 IHC 1:150-1:600 IF/ICC 1:50-1:500

**Applications** 

Tested Applications: IF/ICC, IHC, WB, ELISA Species Specificity: Human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: Caco-2 cells, HeLa cells, LNCaP cells, HEK-293 cells, HepG2 cells, Jurkat cells, HSC-T6 cells, NIH/3T3 cells, Neuro-2a cells

IHC: human breast cancer tissue,

IF/ICC: HepG2 cells,

**Background Information** 

RBM39, also named as HCC1 or RNPC2, is a 530 amino acid protein, which contains three RRM (RNA recognition motif) domains and belongs to the splicing factor SR family. RBM39 is widely expressed in various tissues and with highly expression in pancreas, skeletal muscle, lung and brain. RBM39 is a transcriptional coactivator for steroid nuclear receptors ESR1/ER-alpha and ESR2/ER-beta, and JUN/AP-1. RBM39 may be involved in pre-mRNA splicing process.

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 67420-1-1g (RBM39 antibody) at dilution of 1:2500 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 67420-1-Ig (RBM39 antibody) at dilution of 1:300 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using 67420-1-lg (RBM39 antibody), at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Various lysates were subjected to SDS PAGE followed by western blot with 67420-1-1g (RBM39 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using RBM39 antibody (67420-1-Ig, Clone: 2D2C8) at dilution of 1:2000 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-phalloidin (red).