

For Research Use Only

TAOK3 Monoclonal antibody, PBS Only

Catalog Number: 67451-1-PBS

Featured Product



Basic Information

Catalog Number: 67451-1-PBS	GenBank Accession Number: BC002756	Purification Method: Protein A purification
Size: 1 mg/ml	GeneID (NCBI): 51347	CloneNo.: 1A5A9
Source: Mouse	UNIPROT ID: Q9H2K8	
Isotype: IgG1	Full Name: TAO kinase 3	
Immunogen Catalog Number: AG29146	Calculated MW: 105 kDa	
	Observed MW: 100-105 kDa	

Applications

Tested Applications:
WB, IHC, IF-P, ELISA

Species Specificity:
human, mouse

Background Information

TAOK3 (Serine/threonine-protein kinase TAO3) is also named as DPK, JIK, KDS, MAP3K18 and belongs to the STE Ser/Thr protein kinase family. It acts as an activator of the p38/MAPK14 stress-activated MAPK cascade. In response to DNA damage, it is involved in the G2/M transition DNA damage checkpoint by activating the p38/MAPK14 stress-activated MAPK cascade, probably by mediating phosphorylation of upstream MAP2K3 and MAP2K6 kinases. And this protein can be autophosphorylated (PMID:20949042).

Storage

Storage:
Store at -80°C.
The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:
PBS Only

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67451-1-Ig (TAOK3 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 67451-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded human prostate cancer tissue slide using 67451-1-Ig (TAOK3 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67451-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded human prostate cancer tissue slide using 67451-1-Ig (TAOK3 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67451-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed human prostate cancer tissue using TAOK3 antibody (67451-1-Ig, Clone: 1A5A9) at dilution of 1:400 and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 67451-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed human prostate cancer tissue using TAOK3 antibody (67451-1-Ig, Clone: 1A5A9) at dilution of 1:400 and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 67451-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded mouse kidney tissue slide using 67451-1-Ig (TAOK3 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67451-1-PBS in a different storage buffer formulation.