## For Research Use Only

## ME2 Monoclonal antibody, PBS Only

Catalog Number: 67457-1-PBS

**Featured Product** 



**Purification Method:** 

Protein A purification

CloneNo.:

6E6E2

**Basic Information** 

Catalog Number:

Size: 1 mg/ml

67457-1-PBS

Immunogen Catalog Number:

GeneID (NCBI): 4200

**UNIPROT ID:** Source: Mouse P23368 Isotype: Full Name:

lgG2a malic enzyme 2, NAD(+)-dependent, mitochondrial

AG20741 Calculated MW: 65 kDa

Observed MW: 54 kDa, 65 kDa

BC000147

GenBank Accession Number:

**Applications** 

**Tested Applications:** WB,Indirect ELISA,IHC Species Specificity: Human, Mouse, Rat

## **Background Information**

ME2(malic enzyme 2) is also named as NAD-ME and belongs to the malic enzymes family. ME1, ME2 are important for NADPH production, lipogenesis and glutamine metabolism, but ME2 has a more profound effect. ME2 is a genome-coded mitochondrial enzyme that converts malate to pyruvate and is involved in neuronal synthesis of the neurotransmitter  $\gamma$  -aminobutyric acid (GABA)(PMID:15532013). This protein has 2 isoforms produced by alternative splicing with the MW of 65 kDa and 54 kDa.

Storage


Storage:

Store at -80°C.


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C Storage Buffer:

PBS Only

## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 67457-1-lg (ME2 antibody) at dilution of 1:2500 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 67457-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 67457-1-Ig (ME2 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67457-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 67457-1-Ig (ME2 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67457-1-PBS in a different storage buffer formulation.



WB result of ME2 antibody (67457-1-lg; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-ME2 transfected HeLa cells. This data was developed using the same antibody clone with 67457-1-PBS in a different storage buffer formulation.