For Research Use Only

EEF1A1 Monoclonal antibody

Catalog Number:67495-1-lg 2 Publications




| Basic Information | Catalog Number: 67495-1-lg | GenBank Accession Numbe BC014224 | er: Purification Method: Protein A purification | | |
|------------------------|--|--|---|-------------------------------------|--|
| | Size: 1000 μg/ml | GenelD (NCBI): 1915 | CloneNo.: 1A9F5 | | |
| | Source: Mouse Isotype: IgG1 Immunogen Catalog Number: AG1938 | UNIPROT ID: P68104 | Recommended Dilutions: WB 1:5000-1:50000 | | |
| | | Full Name: eukaryotic translation elor factor 1 alpha 1 | IHC 1:150-1:600 ngation IF 1:50-1:500 | | |
| | | Calculated MW: 462 aa, 50 kDa | | | |
| | | Observed MW: 49 kDa | | | |
| Applications | Tested Applications: | Pos | Positive Controls: | | |
| | FC, IF/ICC, IHC, WB, ELISA Cited Applications: WB, IP | | WB : HeLa cells, HepG2 cells, HEK-293 cells, Jurkat cells, K-562 cells, HSC-T6 cells, PC-12 cells, NIH/3T3 cells, 4T1 cells | | |
| | Species Specificity: Human, mouse, rat | | | pancreas cancer tissue, mouse brain | |
| | Cited Species: human | IF : HepG2 cells, | | | |
| | Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 | | | | |
| Background Information | promotes the GTP-dependent bi a typical housekeeping gene pro | nding of aminoacyl-tRNA to the A oduct required for the maintenand bond against hydrolysis until a c | TP-binding elongation factor family. T -site of ribosomes during protein biosy e of cell proliferation and/or survival. orrect match between the codon on mR | nthesis. It/ In additior | |
| Notable Publications | Author | Pubmed ID Journal | Applic | ation | |
| | Yingjie Hou | 35036957 STAR Proto | c WB,IP | | |
| | Yingjie Hou | 34478637 Cell Chem | Biol WB | | |
| Storage | Storage: Store at -20°C. Stable for one yes Storage Buffer: PBS with 0.02% sodium azide au Aliquoting is unnecessary for -20 | nd 50% glycerol pH 7.3. | | | |


For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67495-1-1g (EEF1A1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




WB result of EEF1A1 antibody (67495-1-lg; 1:20000; incubated at room temperature for 1.5 hours) with sh-Control and sh-EEF1A1 transfected HepG2 cells.



Immunohistochemical analysis of paraffinembedded human pancreas cancer tissue slide using 67495-1-Ig (EEF1A1 antibody) at dilution of 1:300 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using 67495-1-1g (EEF 1A1 antibody), at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



1X10^6 HepG2 cells were intracellularly stained with 0.4 ug Anti-Human EEF1A1 (67495-1-lg, Clone:1A9F5) and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG1 Isotype Control (MOPC-21) (65124-1-lg, Clone: MOPC-21) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).