For Research Use Only

PRDX6 Monoclonal antibody, PBS Only



Catalog Number:67499-1-PBS

Featured Product

Basic Information

Catalog Number: 67499-1-PBS

Size: 1 mg/ml Source: Mouse Isotype: lgG2a

Immunogen Catalog Number:

AG4727

Calculated MW: 224 aa, 25 kDa Observed MW: 25-30 kDa

Purification Method:

Protein A purification

CloneNo.: 3C12D3

Applications

Tested Applications: WB,IP,Indirect ELISA,IF Species Specificity: Human, Pig

Background Information

PRDX6 (Peroxiredoxin-6), also named as AOP2 or KIAA0106, is a unique member of the peroxiredoxin family of antioxidants. PRDX6 is highly expressed in liver and protects cells from oxidative damage by reducing H2O2 and various lipid Peroxides (PMID: 17382207). It can form a dimer(PMID:20500660).PRDX6 is expressed in all major organs, with a particularly high level in lung (PMID:15890616). Prdx6 is detected at approximately 24 to 28 kDa, and can be monosumoylated with the molecular mass of about 40 kDa (PMID: 24910119).

GenBank Accession Number:

BC035857

9588

P30041

GeneID (NCBI):

UNIPROT ID:


Full Name:

peroxiredoxin 6


Storage

Storage: Store at -80°C. Storage Buffer: PBS Only

Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67499-1-lg (PRDX6 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control. This data was developed using the same antibody clone with 67499-1-PBS in a different storage buffer formulation.




SM=10% starting material; UB=10% unbound fraction; IP=immunoprecipitate.


HAP1 lysates prepared and IP of peroxidoxin 6 performed using 2.0 µg of 67499-1-lg coupled to protein G-Sepharose beads. Ponceau stained transfers shown for each blot. Data provided by YCharOS, an open science company with a mission to validate commercial antibodies to improve scientific reproducibility and transparency. This data was developed using the same antibody clone with 67499-1-PBS in a different storage buffer formulation.




HAP1 and HeLa (WT and PRDX6 KO) lysates prepared with RIPA buffer, 25ug protein loaded. 67499-1-lg incubated at 1:1000 at 4°C overnight in 5% BSA in TBST. Ponceau stained transfers shown on right. Data provided by YCharOS, an open science company with a mission to validate commercial antibodies to improve scientific reproducibility and transparency. This data was developed using the same antibody clone with 67499-1-PBS in a different storage buffer formulation.



HAP1 WT cells (yellow outline) and PRDX6 KO cells (red outline) labelled with a green or a far red fluorescence dye, respectively. Cells fixed with 4% PFA and stained with 67499-1-lg at 1:1000 plus DAPI. Bars = 10 μ m. Data provided by YCharOS, an open science company with a mission to validate commercial antibodies to improve scientific reproducibility and transparency. This data was developed using the same antibody clone with 67499-1-PBS in a different storage buffer



WB result of PRDX6 antibody (67499-1-Ig; 1:20000; incubated at room temperature for 1.5 hours) with sh-Control and sh-PRDX6 transfected HeLa cells. This data was developed using the same antibody clone with 67499-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using PRDX6 antibody (67499-1-lg, Clone: 3C12D3) at dilution of 1:800 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-phalloidin (red). This data was developed using the same antibody clone with 67499-1-PBS in a different storage buffer formulation.