For Research Use Only

CDK5R1 Monoclonal antibody

Catalog Number:67503-1-lg 3 Publications



Basic Information

Applications

Catalog Number: 67503-1-lg Size:

lgG2b

1000 µ g/ml
Source:
Mouse
Isotype:

Immunogen Catalog Number:

AG13152

Tested Applications:

IHC, WB, ELISA
Cited Applications:
WB, IF, IHC, CoIP
Species Specificity:

Human, mouse, rat, pig, rabbit

Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

GenBank Accession Number: Purification Method: BC020580 Protein A purification

BC020580 Protein A

Genel D (NCBI): CloneNo.:
8851 1A3A5

 UNIPROT ID:
 Recommended Dilutions:

 Q15078
 WB 1:5000-1:50000

 Full Name:
 IHC 1:200-1:800

cyclin-dependent kinase 5, regulatory

subunit 1 (p35)

Calculated MW:

Observed MW: 34 kDa

34 kDa

Positive Controls:

WB: pig cerebellum tissue, pig brain tissue, rat brain tissue, mouse brain tissue, mouse cerebellum tissue

IHC: mouse brain tissue, rat brain tissue

Background Information

Cyclin dependent kinase-5 (Cdk5), is an atypical but essential member of the Cdk family of proline-directed serine/threonine kinases. CDK5R1 encodes p35, a specific activator of the serine/threonine kinase CDK5, which plays a central role in neuronal migration and differentiation during central nervous system development. The p35 form of this protein is proteolytically cleaved by calpain, generating a p25 form.

Notable Publications

Author	Pubmed ID	Journal	Application
Li Zeng	35024240	Mol Ther Nucleic Acids	IHC
Ran Zhuo	38636893	Cancer Lett	WB
Xiao-Ling Liang	38024857	J Pharm Anal	WB,IF,CoIP

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67503-1-lg (CDK5R1 antibody) at dilution of 1:15000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Alpha Tubulin Monoclonal antibody (HRP-66031) as loading control.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 67503-1-Ig (CDK5R1 antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 67503-1-Ig (CDK5R1 antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



mouse cerebellum tissue were subjected to SDS PAGE followed by western blot with 67503-1-1g (CDK5R1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.