For Research Use Only

TCF12 Monoclonal antibody, PBS Only



Catalog Number: 67523-1-PBS

Basic Information

Catalog Number:

67523-1-PBS

Size: 1mg/ml Source: Mouse Isotype:

lgG2a Immunogen Catalog Number:

AG6287

73 kDa

80 kDa

Observed MW:

GenBank Accession Number:

BC050556

6938

Q99081

Full Name:

transcription factor 12 Calculated MW:

GeneID (NCBI):

UNIPROT ID:

Purification Method:

Protein A purification CloneNo.:

1B8D4

Applications

Tested Applications: Indirect ELISA, WB Species Specificity: rat, mouse, human

Background Information

The product of TCF12 gene (TCF12, also called HTF4 or HEB) is a member of helix-loop-helix (HLH) protein family [PMID:22130667]. TCF 12 is a transcriptional regulator that involved in the initiation of neuronal differentiation. It's also known as E proteins, and activates transcription by binding to the E box (5'-CANNTG-3'), because of their direct DNA (E-box) binding ability [PMID: 19204326]. TCF12 was shown to suppress E-cadherin expression during the early stages of renal tubular epithelial cell dedifferentiation [PMID:17202424].

Storage


Storage:

Store at -80°C.


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer: PBS Only

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67523-1-lg (TCF12 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 67523-1-PBS in a different storage buffer formulation.



1X10^6 HepG2 cells were intracellularly stained with 0.5 ug Anti-Human TCF12 (67523-1-lg, Clone:1B8D4) and Coralite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.5 ug Mouse IgG2b Isotype Control (66360-3-lg, Clone: K11B8C4B5) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011). This data was developed using the same antibody clone with 67523-1-PBS in a different storage buffer