For Research Use Only

CCT8 Monoclonal antibody

Catalog Number:67539-1-lg Featured Product

3 Publications



Basic Information

Catalog Number: 67539-1-lg Size:

1700 µg/ml Source: Mouse Isotype: IgG2a

Immunogen Catalog Number:

AG30082

GenBank Accession Number:

BC012584 GeneID (NCBI): 10694 **UNIPROT ID:** P50990

Full Name: chaperonin containing TCP1, subunit IF 1:750-1:3000 8 (theta)

Calculated MW: 548 aa, 60 kDa

Observed MW: 60 kDa

Applications

Tested Applications: IF/ICC, IHC, WB, ELISA Cited Applications: WB, IP, IHC

Species Specificity: Human, mouse, rat **Cited Species:**

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HEK-293 cells, NIH/3T3 cells, A549 cells, HeLa cells, Jurkat cells, K-562 cells, HSC-T6 cells

Purification Method:

Protein A purification

Recommended Dilutions:

WB 1:5000-1:50000 IHC 1:1000-1:4000

CloneNo.:

1A7C3

IHC: human liver cancer tissue,

IF: MCF-7 cells,

Background Information

CCT8, also named as C21orf112, CCTQ, KIAA0002, CCT-theta, TCP-1-theta and NY-REN-15, belongs to the TCP-1 chaperonin family. It is a molecular chaperone; assists the folding of proteins upon ATP hydrolysis. As part of the BBS/CCT complex, CCT8 may play a role in the assembly of BBSome, a complex involved in ciliogenesis regulating transports vesicles to the cilia. Dysregulation of CCT8 has been linked to proliferation and invasion of tumors.

Notable Publications

Author	Pubmed ID	Journal	Application
Xinyu Ma	35366418	Cell	WB
Zhiqiang Wu	37928427	J Cancer	WB,IHC,IP
Yuichi Yagita	37480851	Cell	WB

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67539-1-1g (CCT8 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 67539-1-1g (CCT8 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 67539-1-1g (CCT8 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Methanol) fixed MCF-7 cells using CCT8 antibody (67539-1-lg, Clone: 1A7C3) at dilution of 1:1500 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



WB result of CCT8 antibody (67539-1-lg; 1:10000; incubated at room temperature for 1.5 hours) with sh-Control and sh-CCT8 transfected HEK-293 cells.