For Research Use Only

Collagen Type V Monoclonal antibody



Catalog Number:67604-1-lg

8 Publications

Basic Information

Catalog Number: 67604-1-lg Size: 1000 µg/ml Source:

Isotype:

lgG1

Mouse

GenBank Accession Number:

NM_000093 GeneID (NCBI): 1289 UNIPROT ID: P20908

Full Name: collagen, type V, alpha 1

Calculated MW: 184 kDa Observed MW: 220 kDa Purification Method:

Protein A purification

CloneNo.: 2C2C2

Recommended Dilutions: WB 1:5000-1:50000 IHC 1:500-1:2000

Applications

Tested Applications: WB, IHC, ELISA Cited Applications: WB, IF, IHC

Species Specificity: Human, Mouse, Rat, Pig Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: HT-1080 cells, mouse colon tissue, HeLa cells, pig colon tissue, rat colon tissue, Saos-2 cells, SiHa cells

IHC: human breast cancer tissue,

Background Information

Notable Publications

Author	Pubmed ID	Journal	Application
Meng-Han Zhang	36273780	Respir Physiol Neurobiol	WB
Zongfu Pan	36192735	Mol Cancer	WB,IHC,IF
Yinan Chen	34976220	Theranostics	IF

Storage

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffe


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




HT-1080 cells were subjected to SDS PAGE followed by western blot with 67604-1-1g (Collagen Type V antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 67604-1-Ig (Collagen Type V antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 67604-1-Ig (Collagen Type V antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 67604-1-1g (Collagen Type V antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 67604-1-Ig (Collagen Type V antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.