

CD24 Monoclonal antibody

Catalog Number: 67627-1-Ig **1 Publications**

Basic Information

Catalog Number:

67627-1-Ig

Size:

1000 µg/ml

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG11679

GenBank Accession Number:

BC007674

GeneID (NCBI):

100133941

UNIPROT ID:

P25063

Full Name:

CD24 molecule

Calculated MW:

8 kDa

Observed MW:

42 kDa

Purification Method:

Protein A purification

CloneNo.:

1H5C4

Recommended Dilutions:

WB 1:5000-1:50000

Applications

Tested Applications:

FC, WB, ELISA

Cited Applications:

WB

Species Specificity:

Human

Cited Species:

human

Positive Controls:

WB : MCF-7 cells, Raji cells, Ramos cells, Daudi cells, Jurkat cells, K-562 cells, HL-60 cells

Background Information

CD24 (known as heat stable antigen) is a small highly glycosylated GPI-linked sialoprotein. It is normally expressed at the surface of most B lymphocytes and differentiating neuroblasts, and it is also up-regulated in a wide variety of cancers. Studies have shown that CD24 functions in the regulation of B-cell apoptosis, leukocyte signal transduction, and leukocyte adhesion. Since it is highly glycosylated, the apparent molecular weight of CD24 could be variable, ranging from 30 kDa to 70 kDa. (Ref: Akihiko Sano, MD., 2009)

Notable Publications

Author	Pubmed ID	Journal	Application
Xudong Zhu	38164371	Breast Cancer (Dove Med Press)	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67627-1-Ig (CD24 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



1X10⁶ Ramos cells were intracellularly stained with 0.4 ug Anti-Human CD24 (67627-1-Ig, Clone:1H5C4) and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).