

For Research Use Only

ITK Monoclonal antibody, PBS Only

Catalog Number: 67703-1-PBS



Basic Information

Catalog Number:

67703-1-PBS

Size:

1 mg/ml

Source:

Mouse

Isotype:

IgG2a

Immunogen Catalog Number:

AG17087

GenBank Accession Number:

BC109077

GeneID (NCBI):

3702

UNIPROT ID:

Q08881

Full Name:

IL2-inducible T-cell kinase

Calculated MW:

620 aa, 72 kDa

Observed MW:

70 kDa

Purification Method:

Protein A purification

CloneNo.:

1F9D8

Applications

Tested Applications:

WB, Indirect ELISA, IF, FC

Species Specificity:

Human

Background Information

ITK is a member of Tec family (BTK, ITK, Tec, BMX and RLK), which expressed in normal T-lymphocytes and T-cell associated hematopoietic malignancies and have confirmed its critical role in regulating T lymphocyte function in EBV-driven lymphoproliferative disease and immune-mediated disorders. Tec kinase family members shares similarities structure, consisting of PH domain, SH3 domain, SH2 domain and kinase domain. Loss of function mutations in ITK results in hypogammaglobulinemia and CD4+ T cell loss in humans, and the patients often present with EBV-associated B cell lymphoproliferative syndrome (PMID: 30814910, PMID: 31025232).

Storage

Storage:

Store at -80°C.

Storage Buffer:

PBS Only

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




PBMC cells were subjected to SDS PAGE followed by western blot with 67703-1-Ig (ITK antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 67703-1-PBS in a different storage buffer formulation.



Jurkat cells were subjected to SDS PAGE followed by western blot with 67703-1-Ig (ITK antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 67703-1-PBS in a different storage buffer formulation.



1X10⁶ Jurkat cells were intracellularly stained with 0.2 ug Anti-Human ITK (67703-1-Ig, Clone:1F9D8) and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (green), and 0.2 ug Mouse IgG2a Isotype Control (66360-2-Ig, Clone: K11A1B2A2) (black). Cells were fixed with 90% MeOH. This data was developed using the same antibody clone with 67703-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed Jurkat cells using ITK antibody (67703-1-Ig, Clone: 1F9D8) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 67703-1-PBS in a different storage buffer formulation.