For Research Use Only

VAMP2 Monoclonal antibody, PBS Only

Size: 1 mg/ml



Purification Method:

Protein A purification

CloneNo.:

1G7E8

Catalog Number: 67822-1-PBS

Basic Information

Catalog Number: GenBank Accession Number: 67822-1-PBS

BC002737 GeneID (NCBI): 6844

UNIPROT ID: Source: Mouse P63027 Isotype: Full Name:

lgG3 vesicle-associated membrane protein

2 (synaptobrevin 2) Immunogen Catalog Number: AG17908 Calculated MW: 13 kDa

> Observed MW: 19 kDa

Applications

Tested Applications: WB,Indirect ELISA,IHC,IF Species Specificity:

Human, Mouse, Rat, Pig, Rabbit


Background Information

VAMP2 (vesicle-associated membrane protein 2), also named as synaptobrevin 2, is a member of the SNARE (soluble NSF-attachment protein receptor) family proteins. Characterized by a common sequence called the SNARE motif, SNARE proteins are involved in membrane fusion and vesicular transport (PMID: 11252968). VAMP2, with a molecular mass of 15-19 kDa, consists of a short N-terminal sequence, a SNARE motif, and a C-terminal transmembrane region. It is required for fast calcium-triggered synaptic vesicle fusion. VAMP2 forms a stable complex with STX1 (syntaxin 1) and SNAP25 (synaptosomal-associated protein 25) during synaptic vesicle fusion $(PMID: 16793874). \ It also forms a distinct complex with synaptophysin. \ VAMP2 is expressed in nervous system and a synaptophysin of the complex with synaptophysin of the complex with synaptophysin. \ VAMP2 is expressed in nervous system and the complex with synaptophysin of the complex with synaptophysin o$ some non-neuronal tissues, such as skeletal muscle (PMID: 18570252).

Storage

Storage: Store at -80°C. Storage Buffer: PBS Only

Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67822-1-lg (VAMP2 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 67822-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 67822-1-Ig (VAMP2 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67822-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 67822-1-Ig (VAMP2 antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67822-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed U-87 MG cells using VAMP2 antibody (67822-1-Ig, Clone: 1G7E8) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-phalloidin (red). This data was developed using the same antibody clone with 67822-1-PBS in a different storage buffer formulation.