

For Research Use Only

C9orf72 Monoclonal antibody

Catalog Number: 67824-1-Ig



Basic Information

Catalog Number: 67824-1-Ig	GenBank Accession Number: BC020851	Purification Method: Protein A purification
Size: 1000 µg/ml	GeneID (NCBI): 203228	CloneNo.: 2F9A2
Source: Mouse	UNIPROT ID: Q96LT7	Recommended Dilutions: WB 1:1000-1:6000
Isotype: IgG2a	Full Name: chromosome 9 open reading frame 72	
Immunogen Catalog Number: AG21080	Calculated MW: 481 aa, 54 kDa	
	Observed MW: 50 kDa	

Applications

Tested Applications: FC, WB, ELISA	Positive Controls: WB : pig brain tissue, HEK-293 cells, U-87 MG cells, Neuro-2a cells, PC-12 cells, C6 cells, rat brain tissue, mouse brain tissue
Species Specificity: Human, mouse, rat, pig	

Background Information

C9ORF72 has a domain with polymorphic hexanucleotide repeat (GGGGCC). The C9ORF72-hexanucleotide repeat expansions have been recently identified as genetic markers in amyotrophic lateral sclerosis (ALS) and frontotemporal lobar degeneration (FTLD). The C9ORF72 repeat expansions may indicate a worse prognosis in ALS. Human C9ORF72 has some isoforms with MW 54-60 kDa and 25-30 kDa. Mouse C9orf72 has some isoforms with MW 50-60 kDa and 35 kDa.

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67824-1-Ig (C9orf72 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.



1×10^6 HEK-293T cells were intracellularly stained with 0.4 μ g Anti-Human C9orf72 (67824-1-Ig, Clone:2F9A2) and CoraLite[®]488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 μ g Mouse IgG2a Isotype Control (C1.18.4) (65208-1-Ig, Clone: C1.18.4) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Neuro-2a cells were subjected to SDS PAGE followed by western blot with 67824-1-Ig (C9orf72 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.