

For Research Use Only

ROR2 Monoclonal antibody

Catalog Number: 67906-1-Ig **1 Publications**



Basic Information

Catalog Number: 67906-1-Ig	GenBank Accession Number: BC130522	Purification Method: Protein G purification
Size: 1000 µg/ml	GeneID (NCBI): 4920	CloneNo.: 1A12C5
Source: Mouse	UNIPROT ID: Q01974	Recommended Dilutions: WB 1:5000-1:50000
Isotype: IgG1	Full Name: receptor tyrosine kinase-like orphan receptor 2	
Immunogen Catalog Number: AG31197	Calculated MW: 943 aa, 105 kDa	
	Observed MW: 105 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : HEK-293 cells, HSC-T6 cells, NIH/3T3 cells, MDA-MB-231 cells, T-47D cells, HeLa cells, K-562 cells, Jurkat cells, 4T1 cells
Cited Applications: WB	
Species Specificity: Human, Mouse, Rat	
Cited Species: mouse	

Background Information

Tyrosine-protein kinase transmembrane receptor ROR2 belongs to the ROR family of receptor tyrosine kinases (RTKs) (PMID: 12815588). ROR2 is a developmentally regulated protein that has significant expression and roles in a wide variety of tissues during early embryonic development. It was established that ROR2 plays a key role in skeletal and neural organogenesis, as well as in midgut development in the mouse, but then becomes repressed in adult tissues. It may act as a receptor for wnt ligand WNT5A which may result in the inhibition of WNT3A-mediated signaling (PMID: 25614331). Mutations in the ROR2 gene are associated with autosomal recessive Robinow syndrome and autosomal dominant brachydactyly type B (PMID: 12815588).

Notable Publications

Author	Pubmed ID	Journal	Application
Yitian Lu	38519978	J Nanobiotechnology	WB

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67906-1-Ig (ROR2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Alpha Tubulin Monoclonal antibody (HRP-66031) as loading control.