For Research Use Only

USP10 Monoclonal antibody, PBS Only



Catalog Number: 67917-1-PBS

Basic Information

Catalog Number:

GenBank Accession Number: BC000263

Purification Method: Protein A purification

67917-1-PBS

GeneID (NCBI):

Size: 1mg/ml Source:

9100 **UNIPROT ID:**

Q14694

CloneNo.: 1B10C9

Mouse Isotype:

Full Name:

lgG1

ubiquitin specific peptidase 10

Calculated MW: Immunogen Catalog Number:

AG7233

87 kDa, 92 kDa Observed MW:

100-110 kDa

Applications

Tested Applications:

WB, IF, IHC, Indirect ELISA

Species Specificity:

Human, mouse, rat

Background Information


USP10, a cytoplasmic ubiquitin-specific protease, deubiquitinates p53, reversing Mdm2-induced p53 nuclear export and degradation. After DNA damage, USP10 is stabilized, and a fraction of USP10 translocates to the nucleus to activate p53. The translocation and stabilization of USP10 is regulated by ATM-mediated phosphorylation of USP10 at Thr42 and Ser337. (PMID: 20096447, PMID: 26876938)

Storage


Storage: Store at -80°C.

Storage Buffer: PBS Only


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67917-1-1g (USP10 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Alpha Tubulin Monoclonal antibody (HRP-66031) as loading control. This data was developed using the same antibody clone with 67917-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 67917-1-lg (USP10 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67917-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HEK-293 cells using USP10 antibody (67917-1-lg, Clone: 1B10C9) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 67917-1-PBS in a different storage buffer formulation.



WB result of USP10 antibody (67917-1-1g; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-USP10 transfected MCF-7 cells. This data was developed using the same antibody clone with 67917-1-PBS in a different storage buffer formulation.