For Research Use Only

IPO5 Monoclonal antibody


Catalog Number:67962-1-Ig




Basic Information	Catalog Number: 67962-1-lg	GenBank Accession Num BC001497	ber: Purification Method: Protein A purification	
	Size: 1000 μg/ml	GeneID (NCBI): 3843	CloneNo.: 1E2B9	
	Source: Mouse	UNIPROT ID: 000410	Recommended Dilutions: WB 1:1000-1:5000	
	Isotype: IgG1 Immunogen Catalog Number: AG30955	Full Name: importin 5	IHC 1:250-1:1000 IF/ICC 1:200-1:800	
		Observed MW: 124 kDa		
Applications	IF/ICC, IHC, WB, ELISA WB : Species Specificity: cells,		ositive Controls:	
			WB : HeLa cells, HEK-293 cells, HepG2 cells, Jurkat cells, K-562 cells, HSC-T6 cells, PC-12 cells, NIH/3T3 cells	
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0		IHC : mouse testis tissue,	
			/ICC : A431 cells,	
Background Information	IPO5 is a member of the importin beta family. Structurally, the protein adopts the shape of a right-hand solenoid and is composed of 24 HEAT repeats. IPO5 facilitates cytoplasmic polyadenylation element-binding protein CPEB3 translocation by binding to the RRM1 motif of CPEB3 in neurons. NMDAR signaling increases RanBP1 expression and reduces the level of cytoplasmic GTP-bound Ran. These changes enhance CPEB3-IPO5 interaction, which consequently accelerates the nuclear import of CPEB3 and promotes its nuclear function.			
Storage	Storage: Store at -20°C. Stable for one year Storage Buffer: PBS with 0.02% sodium azide and Aliquoting is unnecessary for -20°	I 50% glycerol pH 7.3.		

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67962-1-1g (IPO5 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with Lamin B1 Monoclonal antibody (66095-1-1g) as loading control.



Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 67962-1-Ig (IPO5 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Methanol) fixed A431 cells using IPO5 antibody (67962-1-lg, Clone: 1E2B9) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).