For Research Use Only

MYBBP1A Monoclonal antibody

Catalog Number:67996-1-Ig




Basic Information	Catalog Number: GenBank Accession Numb 67996-1-lg BC050546		lumber:	Purification Method: Protein G purification
	Size: GeneID (NCBI): 1000 μ g/ml 10514			CloneNo.: 1H2D10
	Source:UNIPROT ID:MouseQ9BQGO			Recommended Dilutions: WB 1:5000-1:50000 IHC 1:500-1:2000 IF/ICC 1:300-1:1200
	lsotype: lgG1	Full Name: MYB binding protein (P160) 1a Calculated MW: 149 kDa		
	Immunogen Catalog Number: AG6008			
		Observed MW: 140 kDa		
Applications	Tested Applications: IF/ICC, IHC, WB, ELISA Species Specificity: Human, mouse, rat		Positive Controls: WB : LNCaP cells, HSC-T6 cells, NIH/3T3 cells, 4T1 cells, HeLa cells, HEK-293 cells, HepG2 cells, Jurkat cells, K-562 cells	
	IF/ICC : HepG2 cells, NIH/3T3 cells			
	Background Information	The protooncogene MYB is predominantly expressed in immature hemopoietic cells where it has an essential role i hemopoietic cell proliferation and differentiation. Oncogenically activated forms of MYB is generally N- and/or C- terminal truncations of the normal MYB protein. Removal of the C terminus of MYB disrupts or deletes a region termed the negative regulatory domain (NRD), resulting in an increase in DNA binding, transactivation, and transformation by MYB. One feature of the NRD is a leucine zipper-like motif [PMID: 8302594]. Murine Myb-binding protein-1a (MYBBP1A), originally called P160, was identified by its ability to interact specifically with Myb via this leucine zipper-like motif. MYBBP1A modulates MYB activity upon binding to the MYB NRD [PMID: 10644447, 9447996].		
Storage	Storage: Store at -20°C. Storage Buffer: PBS with 0.02% sodium azide and	50% glycerol nH 7 3		

 For technical support and original validation data for this product please contact:

 T: 4006900926
 E: Proteintech-CN@ptglab.com
 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67996-1-1g (MYBBP1A antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




HSC-T6 cells were subjected to SDS PAGE followed by western blot with 67996-1-1g (MYBBP1A antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 67996-1-Ig (MYBBP1A antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using MYBBP1A antibody (67996-1-lg, Clone: 1H2D10) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-Phalloidin (red).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using MYBBP1A antibody (67996-1-lg, Clone: 1H2D10) at dilution of 1:600 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-Phalloidin (red).



Immunofluorescent analysis of (4% PFA) fixed NIH/3T3 cells using MYBBP1A antibody (67996-1-Ig, Clone: 1H2D10) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed NIH/3T3 cells using MYBBP1A antibody (67996-1-Ig, Clone: 1H2D10) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).