For Research Use Only

## MYBBP1A Monoclonal antibody, PBS Only



Catalog Number: 67996-1-PBS

**Basic Information** 

Catalog Number:

BC050546

Purification Method:

67996-1-PBS

GeneID (NCBI):

Protein G purification

Size: 1 mg/ml

10514

CloneNo.: 1H2D10

1 mg/m Source:

UNIPROT ID: Q9BQG0 Full Name:

Mouse Isotype: IgG1

AG6008

MYB binding protein (P160) 1a

GenBank Accession Number:

Immunogen Catalog Number:

Calculated MW: 149 kDa

149

Observed MW:

140 kDa

Applications

**Tested Applications:** 

WB,Indirect ELISA,IHC,IF

Species Specificity:

Human, mouse, rat

## **Background Information**

The protooncogene MYB is predominantly expressed in immature hemopoietic cells where it has an essential role in hemopoietic cell proliferation and differentiation. Oncogenically activated forms of MYB is generally N- and/or C-terminal truncations of the normal MYB protein. Removal of the C terminus of MYB disrupts or deletes a region termed the negative regulatory domain (NRD), resulting in an increase in DNA binding, transactivation, and transformation by MYB. One feature of the NRD is a leucine zipper-like motif [PMID: 8302594]. Murine Myb-binding protein-1a (MYBBP1A), originally called P160, was identified by its ability to interact specifically with Myb via this leucine zipper-like motif. MYBBP1A modulates MYB activity upon binding to the MYB NRD [PMID: 10644447, 9447996].

Storage


Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer: PBS Only


## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67996-1-lg (MYBBP1A antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 67996-1-PBS in a different storage buffer formulation




HSC-T6 cells were subjected to SDS PAGE followed by western blot with 67996-1-lg (MYBBP1A antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 67996-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 67996-1-lg (MYBBP1A antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67996-1-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using MYBBP1A antibody (67996-1-lg, Clone: 1H2D10) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-Phalloidin (red). This data was developed using the same antibody clone with 67996-1-PBS in a different storage buffer formulation



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using MYBBP1A antibody (67996-1-lg, Clone: 1H2D10) at dilution of 1:600 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-Phalloidin (red). This data was developed using the same antibody clone with 67996-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed NIH/3T3 cells using MYBBP1A antibody (67996-1-lg, Clone: 1H2D10) at dilution of 1:400 and Coralite® 488-Conjugated AffiniPure Goat Anti-Mouse IgC(H+I). This data was developed using the same antibody clone with 67996-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed NIH/3T3 cells using MYBBP1A antibody (67996-1-Ig, Clone: 1H2D10) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 67996-1-PBS in a different storage buffer formulation.