For Research Use Only

PGK1 Monoclonal antibody

Catalog Number:68035-1-lg 1 Publications



Basic Information

Catalog Number: 68035-1-lg

Size: 1000 μ g/ml

Source: Mouse

Isotype:

lgG1

GenBank Accession Number: BC103752 GeneID (NCBI): 5230 UNIPROT ID: P00558

Full Name: I phosphoglycerate kinase 1

Immunogen Catalog Number:Calculated MW:AG12099417 aa, 45 kDa

Observed MW: 40-45 kDa Purification Method:

1B2F3

Protein G purification CloneNo.:

Recommended Dilutions: WB 1:5000-1:50000 IHC 1:1000-1:4000

Applications

Tested Applications: IHC, WB, ELISA Cited Applications: WB

Species Specificity: Human, Mouse, Rat Cited Species: human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: A431 cells, NCCIT cells, PC-3 cells, LNCaP cells, HeLa cells, Jurkat cells, human platelet tissue, rat testis tissue, mouse testis tissue

IHC: human liver cancer tissue, mouse kidney tissue, rat kidney tissue, rat testis tissue

Background Information

PGK1, also known as PGKA, belongs to the phosphoglycerate kinase family. PGK1 is the first ATP-generating enzyme in glycolysis, which catalyzes the conversion of 1,3-diphosphoglycerate (1,3-BPG) to 3- phosphoglycerate (3-PG) and produces one molecule of ATP. PGK1 has other functions besides regulating glycolytic metabolism, including mediating autophagy initiation, DNA replication and repair in mammal cell nuclei. PGK1 has 2 isoforms with the molecular mass of 41 and 45 kDa (PMID: 31815035, PMID: 31911580).

Notable Publications

Author	Pubmed ID	Journal	Application
Wenke Bai	38043497	Transl Oncol	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



A431 cells were subjected to SDS PAGE followed by western blot with 68035-1-lg (PGK1 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 68035-1-Ig (PGK1 antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 68035-1-Ig (PGK1 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).