#### For Research Use Only

# BNIP3 Monoclonal antibody

Catalog Number: 68091-1-lg 17 Publications



**Basic Information** 

Catalog Number: 68091-1-lg Size:

1000 µg/ml Source: Mouse Isotype: lgG2b

Immunogen Catalog Number:

AG27935

GenBank Accession Number:

BC009342 GeneID (NCBI):

Q12983 Full Name: BCL2/adenovirus E1B 19kDa

**UNIPROT ID:** 

interacting protein 3 Calculated MW: 194 aa, 22 kDa

Observed MW: 25-30,50-60 kDa **Purification Method:** 

Protein A purification

CloneNo.: 2H7D8

Recommended Dilutions: WB 1:5000-1:50000 IF 1:200-1:800

**Applications** 

**Tested Applications:** IF/ICC, WB, ELISA Cited Applications: WB, IF, IHC

Species Specificity: Human, Mouse, rat **Cited Species:** human, rat, mouse

Positive Controls:

WB: HepG2 cells, HSC-T6 cells, NIH/3T3 cells, LNCaP

IF: Cobalt Chloride treated HepG2 cells,

## **Background Information**

BNIP3 (BCL2 interacting protein 3) is a member of the Bcl-2 family that is expressed in mitochondria and induces apoptosis (PMID: 10891486). It can regulate the permeabillity state of outer mitchondrial member (OMM), this regulation is accompanied by the formation of homo- and hetero-oligomers inside the membrane. Hypoxia can lead to transcriptional activation of BNIP3 through HIF-1 a (PMID: 11559532). It's calculated molecule weight is 21.5 kDa. The antibody can detect bands around 25 and 50 kDa, representing monomeric and dimeric forms, respectively (PMID: 21364626).BNIP3 is related to the occurrence and development of tumors, cardiovascular diseases, and Parkinson's diseases.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Xudong Xie	36284303	Cell Commun Signal	WB,IHC
Wenshuo Zhang	36578780	Front Cell Dev Biol	WB
Mei-Fei Wu	36476627	Cell Biosci	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

### **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 68091-1-1g (BNIP3 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed Cobalt Chloride treated HepG2 cells using BNIP3 antibody (68091-1-Ig, Clone: 2H7D8) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).