For Research Use Only

## PACSIN1 Monoclonal antibody, PBS Only



**Purification Method:** 

Protein G purification

CloneNo.:

1B7B4

Catalog Number: 68115-1-PBS

**Basic Information** 

Catalog Number: GenBank Accession Number: 68115-1-PBS BC040228

68115-1-PBS BC040228
Size: GeneID (NCBI):

1 mg/ml29993Source:UNIPROT ID:MouseQ9BY11Isotype:Full Name:

IgG1 protein kinase C and casein kinase

Immunogen Catalog Number: substrate in neurons 1

AG4102 Calculated MW:

444 aa, 51 kDa Observed MW: 48-51 kDa

**Applications** 

Tested Applications: WB,Indirect ELISA,IHC,IF,FC

Species Specificity:

Human, Mouse, Rat, Rabbit, Chicken, Pig


**Background Information** 

PACSIN1 (also known as syndapin-1) is a member of the protein kinase C and casein kinase substrate in neurons (PACSIN) family. In mammals, the PACSIN family is comprised of three members, PACSIN1, PACSIN2, and PACSIN3 (PMID: 34990060). PACSIN1 is expressed mainly in neurons, whereas PACSIN2 is ubiquitously expressed in all tissues, and PACSIN3 is expressed mainly in skeletal muscle and the heart (PMID: 23668323). All of these three members contain an N-terminal F-BAR domain and a C-terminal SH3 domain. PACSIN1 plays a role in endocytosis and endosomal recycling. Meanwhile, it has a role in actin remodeling and microtubule nucleation and also plays a particular role in membrane shaping and reconstruction (PMID: 23035120; 34422904). PACSIN1 is involved in neuromorphogenesis and the regulation of the nervous system, and the inappropriate expression of PACSIN1 has been associated with some neurological diseases, including schizophrenia, Alzheimer's disease, and Huntington's disease (PMID: 34990060).


Storage

Storage: Store at -80°C. Storage Buffer: PBS Only


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 68115-1-lg (PACSIN1 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control. This data was developed using the same antibody clone with 68115-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded mouse cerebellum tissue slide using 68115-1-1g (PACSIN1 antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 68115-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded mouse cerebellum tissue slide using 68115-1-lg (PACSIN1 antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 68115-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Ethanol) fixed SH-SY5Y cells using PACSIN1 antibody (68115-1-Ig, Clone: 1B7B4) at dilution of 1:800 and CoraLite® 488-Conjugated Affini Pure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 68115-1-PBS in a different storage buffer formulation.



1X10^6 SH-SY5Y cells were intracellularly stained with 0.4 ug Anti-Human PACSIN1 (68115-1-lg, Clone:18784) and Coralite® 488-Conjugated AffiniPure Goat Anti-Mouse IgC(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgC1 Isotype Control (MOPC-21) (65124-1-lg, Clone: MOPC-21) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 68115-1-PBS in a

