

For Research Use Only

DOHH Monoclonal antibody, PBS Only



Catalog Number: 68151-1-PBS

Basic Information

Catalog Number: 68151-1-PBS	GenBank Accession Number: BC002817	Purification Method: Protein G purification
Size: 1mg/ml	GeneID (NCBI): 83475	CloneNo.: 1H5H7
Source: Mouse	UNIPROT ID: Q9BU89	
Isotype: IgG1	Full Name: deoxyhypusine hydroxylase/monooxygenase	
Immunogen Catalog Number: AG13041	Calculated MW: 33 kDa	
	Observed MW: 33 kDa	

Applications

Tested Applications:
WB, Indirect ELISA

Species Specificity:
Human, Rat, Mouse

Background Information

Deoxyhypusine hydroxylase (DOHH) is the enzyme catalyzing the second step in the post-translational synthesis of hypusine [N ϵ -(4-amino-2-hydroxybutyl)lysine] in the eukaryotic initiation factor 5A (eIF5A). DOHH has been reported to mediate several crucial cellular functions, including cellular proliferation, differentiation and apoptosis. Moreover, previous studies have established that DOHH is highly involved in several essential biological processes driving human diseases including cancer growth, malarial drug resistance, and HIV-1 replication. In particular, the DOHH/eIF5A signaling pathway was revealed to mediate the ability of nerve growth factor to enhance neuronal growth and survival, highlighting the importance of DOHH in neuroprotection (PMID: 35007708, PMID: 16371467, PMID: 22908221, PMID: 19706422).

Storage

Storage:
Store at -80°C.
The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:
PBS Only

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68151-1-Ig (DOHH antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Alpha Tubulin Monoclonal antibody (HRP-66031) as loading control. This data was developed using the same antibody clone with 68151-1-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 68151-1-Ig (DOHH antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 68151-1-PBS in a different storage buffer formulation.