For Research Use Only

EIF3B Monoclonal antibody, PBS Only

Catalog Number: 68202-1-PBS



Purification Method:

Protein G purification

CloneNo.:

1G12G12

Basic Information

Catalog Number:

68202-1-PBS

Size: 1 mg/ml Source:

Mouse Isotype: lgG1

Immunogen Catalog Number:

AG31427

GenBank Accession Number:

BC001173 GeneID (NCBI):

UNIPROT ID: P55884

Full Name:

93 kDa

8662

eukaryotic translation initiation

factor 3, subunit B Calculated MW:

Observed MW: 115 kDa

Applications

Tested Applications:

WB,Indirect ELISA

Species Specificity: Human, mouse, rat

Background Information

EIF3B, also named Eukaryotic translation initiation factor 3 subunit B, is an 814 amino acid protein, which contains 1 RRM (RNA recognition motif) domain and 8 WD repeats and belongs to the eIF-3 subunit B family. EIF3B as an RNAbinding component of the eukaryotic translation initiation factor 3 (eIF-3) complex, is required for several steps in the initiation of protein synthesis. The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNAi and eIF-5 to form the 43S pre-initiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The $eIF-3\ complex\ is\ also\ required\ for\ disassembly\ and\ recycling\ of\ post-termination\ ribosomal\ complexes\ and$ subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation. The eIF-3 complex specifically targets and initiates translation of a subset of mRNAs involved in cell proliferation, including cell cycling, differentiation, and apoptosis, and uses different modes of RNA stem-loop binding to exert either translational activation or repression. The calculated molecular weight of EIF3B is 93 kDa, but the phosphorylated EIF3B protein is about 115 kDa.

Storage

Storage:


Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:

PBS Only

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68202-1-1g (EIF3B antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Lamin B1 Monoclonal antibody (HRP-66095) as loading control. This data was developed using the same antibody clone with 68202-1-PBS in a different storage buffer formulation.