

For Research Use Only

ARF4 Monoclonal antibody, PBS Only



Catalog Number: 68204-1-PBS

Featured Product

Basic Information

Catalog Number:

68204-1-PBS

Size:

1mg/ml

Source:

Mouse

Isotype:

IgG2b

Immunogen Catalog Number:

AG25522

GenBank Accession Number:

BC003364

GeneID (NCBI):

378

UNIPROT ID:

P18085

Full Name:

ADP-ribosylation factor 4

Calculated MW:

180 aa, 21 kDa

Observed MW:

18 kDa

Purification Method:

Protein A purification

CloneNo.:

2A4B8

Applications

Tested Applications:

WB, IF/ICC, Indirect ELISA

Species Specificity:

human, mouse, rat

Background Information

ARF4, also named as ARF2, belongs to the small GTPase superfamily and Arf family. ADP-ribosylation factors (ARFs) are members of the ARF family of GTP-binding proteins of the Ras superfamily, with 20 kDa protein size. ARFs bind and regulate GTP/GDP cycle by alternating between the active GTP-bound and inactive GDP-bound conformations. ARF family proteins are essential and ubiquitous in eukaryotes. Six highly conserved members of the family have been identified in mammalian cells. They function in vesicular traffic and actin remodelling and other bioprocesses in cells. ARF4 and ARF5 are class II ARFs. Their function are not fully described. (PMID: 7759471, PMID: 16042562). This antibody can bind ARFs for the close sequences.

Storage

Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:

PBS Only

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 68204-1-Ig (ARF4 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 68204-1-PBS in a different storage buffer formulation.




A549 cells were subjected to SDS PAGE followed by western blot with 68204-1-Ig (ARF4 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 68204-1-PBS in a different storage buffer formulation.




Various lysates were subjected to SDS PAGE followed by western blot with 68204-1-Ig (ARF4 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 68204-1-PBS in a different storage buffer formulation.




WB result of ARF4 antibody (68204-1-Ig; 1:10000; incubated at room temperature for 1.5 hours) with sh-Control and sh-ARF4 transfected HeLa cells. This data was developed using the same antibody clone with 68204-1-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 68204-1-Ig (ARF4 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 68204-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using ARF4 antibody (68204-1-Ig, Clone: 2A4B8) at dilution of 1:800 and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 68204-1-PBS in a different storage buffer formulation.



1X10⁶ K-562 cells were intracellularly stained with 0.4 ug Anti-Human ARF4 (68204-1-Ig, Clone:2A4B8) and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG2b Isotype Control (66360-3-Ig, Clone: K11B8C4B5) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 68204-1-PBS in a different storage buffer