For Research Use Only

CD59 Monoclonal antibody

Catalog Number: 68222-1-Ig



Basic Information

Catalog Number: 68222-1-Ig Size:

1000 µ g/ml
Source:
Mouse
Isotype:
IgG2b

Immunogen Catalog Number:

AG32834

GenBank Accession Number:

BC001506 GeneID (NCBI): 966 UNIPROT ID: P13987

CD59 molecule, complement regulatory protein

Calculated MW: 19 kDa Observed MW:

Full Name:

18-25 kDa

Purification Method:

Protein A purification CloneNo.:

1E3A1

Recommended Dilutions: WB 1:5000-1:50000 IHC 1:500-1:2000 IF 1:200-1:800

Applications

Tested Applications: WB, IF/ICC,IF-P, IHC, ELISA Species Specificity:

Human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: human colostrum, A549 cells, HeLa cells, MCF-7 cells, HepG2 cells, K-562 cells, Jurkat cells, human placenta tissue

IHC : human placenta tissue, human lung cancer tissue, human urothelial carcinoma tissue

IF: human placenta tissue, BxPC-3 cells

Background Information

CD59, also named as MIC11, MIN1, MIN2, MIN3, MSK21, MIRL, MACIF, HRF20 and 1F5 antigen, is a cell surface molecule glycoprotein with MW 18-25 kDa. It acts as a determinant of proximal-distal cell identity. CD59 acts by binding to the C8 and/or C9 complements of the assembling membrane attack complex (MAC), thereby preventing incorporation of the multiple copies of C9 required for complete formation of the osmolytic pore. It is involved in signal transduction for T-cell activation complexed to a protein tyrosine kinase.

Storage


Storage:

Store at -20°C. Stable for one year after shipment. Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 68222-1-lg (CD59 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human placenta tissue slide using 68222-1-Ig (CD59 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (4% PFA) fixed human placenta tissue using CD59 antibody (68222-1-lg, Clone: 1E3A1) at dilution of 1:400 and CoraLite®594-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunohistochemical analysis of paraffinembedded human placenta tissue slide using 68222-1-Ig (CD59 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed human placenta tissue using CD59 antibody (68222-1-Ig, Clone: 1E3A1) at dilution of 1:400 and CoraLite®594-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Various lysates were subjected to SDS PAGE followed by western blot with 68222-1-1g (CD59 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.