For Research Use Only

PECR Monoclonal antibody, PBS Only **proteintech**®

Catalog Number:68248-1-PBS

Basic Information	Catalog Number: 68248-1-PBS	GenBank Accession Number: BC002529	Purification Method: Protein G purification	
	Size: 1 mg/ml	GeneID (NCBI): 55825	CloneNo.: 2D3G1	
	Source: Mouse	UNIPROT ID: Q9BY49		
	Isotype: IgG1 Immunogen Catalog Number:	Full Name: peroxisomal trans-2-enoyl-CoA reductase		
	AG7022	Calculated MW: 33 kDa		
		Observed MW: 33 kDa		
Applications	Tested Applications: WB,Indirect ELISA			
	Species Specificity: Human, Mouse, Rat			
Background Information	with 86627 bases. It can participa reaction of four long chain fatty ac very long chain fatty acid (VLCFA acyl coenzyme A, which can furthe a peroxisome protein involved in	Peroxisomal trans-2-enoyl-CoA reductase (PECR) is also called TERP and TECR, which is located on chromosome q35 with 86627 bases. It can participate in carbon chain elongation in fatty acid metabolism and catalyze the last reaction of four long chain fatty acid elongation cycles. Each cycle of PECR adds two carbons to the long chain and very long chain fatty acid (VLCFA) chains, reducing the intermediate of trans-2,3-enoyl coenzyme A fatty acid to acyl coenzyme A, which can further extend the carbon chain by entering a new elongation cycle. Meanwhile, PECR is a peroxisome protein involved in fatty acid synthesis and plays an important role in milk fat synthesis. The molecular mass of PECR is 33 kDa. (PMID: 31467878)		
Storage	Storage: Store at -80°C. The product is shipped with ice pa Storage Buffer: PBS Only	Store at -80°C. The product is shipped with ice packs. Upon receipt, store it immediately at -80°C Storage Buffer:		


For technical support and original validation data for this product please contact: T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Antibodies | ELISA kits | Proteins

www.ptglab.com

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68248-1-Ig (PECR antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Beta Actin Monoclonal antibody (HRP-66009) as loading control. This data was developed using the same antibody clone with 68248-1-PBS in a different storage buffer formulation.