

For Research Use Only

MORC3 Monoclonal antibody, PBS Only



Catalog Number: 68370-1-PBS

Basic Information

Catalog Number: 68370-1-PBS	GenBank Accession Number: BC132731	Purification Method: Protein G purification
Size: 1 mg/ml	GeneID (NCBI): 23515	CloneNo.: 1E1E1
Source: Mouse	UNIPROT ID: Q14149	
Isotype: IgG1	Full Name: MORC family CW-type zinc finger 3	
Immunogen Catalog Number: AG21867	Calculated MW: 939 aa, 107 kDa	
	Observed MW: 130 kDa	

Applications

Tested Applications:
WB, Indirect ELISA

Species Specificity:
Human, mouse

Background Information

MORC3, also named as MORC family CW-type zinc finger protein 3 or NXP2, is a 939 amino acid protein, which contains 1 CW-type zinc finger. MORC3 localizes in the nucleus and are expressed in heart, placenta, skeletal muscle, brain, pancreas, lung, liver, but not kidney. MORC3 as Nuclear factor which forms MORC3-NBs (nuclear bodies) via an ATP-dependent mechanism. Sumoylated MORC3-NBs can also associate with PML-NBs and recruits TP53 and SP100 to PML-NBs, thus regulating TP53 activity. It may be required for influenza A transcription during viral infection. The calculated molecular weight of MORC3 is a 107 kDa, but the Sumoylated MORC3 protein is about 130 kDa.

Storage

Storage:
Store at -80°C.
The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:
PBS Only

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68370-1-Ig (MORC3 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Lamin B1 Monoclonal antibody (HRP-66095) as loading control. This data was developed using the same antibody clone with 68370-1-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 68370-1-Ig (MORC3 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Lamin B1 Monoclonal antibody (HRP-66095) as loading control. This data was developed using the same antibody clone with 68370-1-PBS in a different storage buffer formulation.