

For Research Use Only

Histone H2B Monoclonal antibody

Catalog Number: 68393-1-Ig **1 Publications**



Basic Information

Catalog Number: 68393-1-Ig	GenBank Accession Number: BC005827	Purification Method: Protein G purification
Size: 1000 µg/ml	GeneID (NCBI): 8349	CloneNo.: 1E12G3
Source: Mouse	UNIPROT ID: Q16778	Recommended Dilutions: WB 1:5000-1:50000 IHC 1:250-1:1000 IF 1:400-1:1600
Isotype: IgG1	Full Name: histone cluster 2, H2be	
	Calculated MW: 14 kDa	
	Observed MW: 14-17 kDa	

Applications

Tested Applications: FC, IF/ICC, IHC, WB, ELISA	Positive Controls:
Cited Applications: WB	WB: LNCaP cells, HeLa cells, HEK-293 cells, Jurkat cells, HSC-T6 cells, NIH/3T3 cells, chicken cerebellum tissue, zebrafish tissue
Species Specificity: Human, Mouse, Rat, Zebrafish, Chicken	IHC: human lung cancer tissue,
Cited Species: human	IF: HeLa cells,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Histones are nuclear proteins that are classified into five major protein groups: histones H2A, H2B, H3, and H4 are known as the core histones. Post-translationally modified H2B proteins can modulate the nucleosome/chromatin structure or DNA accessibility to affect the transcriptional pathways linked to embryonic development and cell differentiation. Monoubiquitination of histone H2B has emerged as an important chromatin modification with roles not only in transcription but also in cell differentiation, DNA repair or mRNA processing (PMID: 25027370).

Notable Publications

Author	Pubmed ID	Journal	Application
Xiangjun Yin	37919786	Clin Epigenetics	WB


Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:
T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 68393-1-Ig (Histone H2B antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human lung cancer tissue slide using 68393-1-Ig (Histone H2B antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human lung cancer tissue slide using 68393-1-Ig (Histone H2B antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using Histone H2B antibody (68393-1-Ig, Clone: 1E12G3) at dilution of 1:800 and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-Phalloidin (red).



1X10⁶ HeLa cells were intracellularly stained with 0.4 ug Anti-Human Histone H2B (68393-1-Ig, Clone:1E12G3) and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG1 Isotype Control (MOPC-21) (65124-1-Ig, Clone: MOPC-21) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).