For Research Use Only

Histone H2B Monoclonal antibody, PBS Only



Catalog Number: 68393-1-PBS

Basic Information

Catalog Number:

68393-1-PBS

Size: 1 mg/ml Source:

Mouse Isotype:

lgG1

GenBank Accession Number:

BC005827 GeneID (NCBI):

UNIPROT ID: Q16778 Full Name:

histone cluster 2, H2be

Calculated MW: 14 kDa Observed MW: 14-17 kDa Purification Method: Protein G purification

CloneNo.: 1E12G3

Applications

Tested Applications:

WB, IF, IHC, Indirect ELISA

Species Specificity:

Human, Mouse, Rat, Zebrafish, Chicken


Background Information

Histones are nuclear proteins that are classified into five major protein groups: histones H2A, H2B, H3, and H4 are known as the core histones. Post-translationally modified H2B proteins can modulate the nucleosome/chromatin structure or DNA accessibility to affect the transcriptional pathways linked to embryonic development and cell differentiation. Monoubiquitination of histone H2B has emerged as an important chromatin modification with roles not only in transcription but also in cell differentiation, DNA repair or mRNA processing(PMID: 25027370).


Storage

Storage: Store at -80°C. Storage Buffer: PBS Only


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68393-1-1g (Histone H2B antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 68393-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 68393-1-1g (Histone H2B antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 68393-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 68393-1-1g (Histone H2B antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 68393-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using Histone H2B antibody (68393-1-lg, Clone: 1E12G3) at dilution of 1:800 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-Phalloidin (red). This data was developed using the same antibody clone with 68393-1-PBS in a different storage buffer formulation.