## For Research Use Only

## ATPBD4 Monoclonal antibody

Catalog Number: 68397-1-Ig



**Basic Information** 

Catalog Number:

GenBank Accession Number: BC008485

Purification Method: Protein G purification

68397-1-lg

\_\_\_\_\_

rotein a purincatio

Size: 1000 μg/ml GeneID (NCBI): 89978 CloneNo.: 2B12E1

Source:

UNIPROT ID: Q7L8W6 Recommended Dilutions: WB 1:5000-1:50000

Mouse Isotype:

AG33218

Full Name:

lgG1

ATP binding domain 4

Immunogen Catalog Number:

Calculated MW:

267 aa, 30 kDa

Observed MW:

28-30 kDa

**Applications** 

**Tested Applications:** 

Positive Controls:

WB, ELISA

WB: A549 cells, K-562 cells, LNCaP cells, HeLa cells,

Species Specificity:

HEK-293 cells

Human

**Background Information** 

Diphthamide is a highly modified histidine in the transcription factor EEF2 that is conserved from yeast to human. ATPBD4, also known as DPH6, belongs to the Diphthine--ammonia ligase family. DPH6 amidase may catalyze the last step of diphthamide biosynthesis using ammonium and ATP to play a role in catalyzing the final step of diphthamide biosynthesis, amidation of diphthine to diphthamide (PubMed:23169644).

Storage

Storage:


Store at -20°C. Stable for one year after shipment.


Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

## **Selected Validation Data**





Various lysates were subjected to SDS PAGE followed by western blot with 68397-1-1g (ATPBD4 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Alpha Tubulin Monoclonal antibody (HRP-66031) as loading control.

K-562 cells were subjected to SDS PAGE followed by western blot with 68397-1-1g (ATPBD4 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.