

For Research Use Only

DIABLO Monoclonal antibody

Catalog Number: 68480-1-Ig



Basic Information

Catalog Number: 68480-1-Ig	GenBank Accession Number: BC011909	Purification Method: Protein G purification
Size: 500 µg/ml	GeneID (NCBI): 56616	CloneNo.: 2F12C9
Source: Mouse	UNIPROT ID: Q9NR28	Recommended Dilutions: WB 1:5000-1:50000
Isotype: IgG1	Full Name: diablo homolog (Drosophila)	
Immunogen Catalog Number: AG32774	Calculated MW: 27 kDa	
	Observed MW: 20 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : U2OS cells, K-562 cells, LNCaP cells, HeLa cells, HEK-293 cells, Jurkat cells, HSC-T6 cells
Species Specificity: Human, rat	

Background Information

Smac/DIABLO is a mitochondrial protein that promotes apoptosis by neutralizing members of the IAP family. SMAC/DIABLO is ubiquitously expressed in normal tissues, and the expression of SMAC messenger RNA (mRNA) in adult testis is the highest. Smac/DIABLO encodes a 239 amino acid protein (~27 kDa). The first 55 amino acids are a mitochondrial targeting signal peptide that is cleaved to generate mature Smac-DIABLO (~21 kDa). When released from mitochondria, Smac-DIABLO acts as a natural antagonist of inhibitor of apoptosis proteins (IAPs). Smac-DIABLO antagonistic activity is based on its N-terminal tetrapeptide (AVPI) that binds baculoviral IAP repeat (BIR) domains of IAPs, releasing their inhibitory effects on both initiator and effector caspases, thus promoting cell death. (PMID: 32325691, PMID: 32575872, PMID: 25650938)

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68480-1-Ig (DIABLO antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.



Various lysates were subjected to SDS PAGE followed by western blot with 68480-1-Ig (DIABLO antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.