For Research Use Only

## Histone H3 (C-terminal) Monoclonal antibody, PBS Only



**Purification Method:** 

Protein A purification

CloneNo.:

2G10C3

Catalog Number: 68503-1-PBS

**Basic Information** 

Catalog Number:

68503-1-PBS

Size: 1 mg/ml Source:

Mouse Isotype:

lgG2a

GenBank Accession Number:

BC066245

GeneID (NCBI): 8350

UNIPROT ID: P68431

Full Name:

histone cluster 1, H3a

Observed MW: 15 kDa

**Applications** 

Tested Applications:

WB,Indirect ELISA,IHC

Species Specificity:

Human, mouse, rat, chicken, zebrafish, wheat

**Background Information** 


Histones are small, highly basic proteins that consist of a globular domain with unstructured N- and C-terminal tails protruding from the main structure. Histone H3 is one of the five main histones that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. In addition to their role in DNA compartmentalization, histones also play crucial roles in various biologic processes, including gene expression and regulation, DNA repair, chromatin condensation, cell cycle progression, chromosome segregation, and apoptosis. The ability of histones to regulate chromatin dynamics primarily originates from various posttranslational modifications carried out by histone-modifying enzymes.

Storage

Storage:

Store at -80°C. Storage Buffer: PBS Only


## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 68503-1-lg (HIST1H3A antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 68503-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 68503-1-Ig (HIST1H3A antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 68503-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 68503-1-Ig (HIST1H3A antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 68503-1-PBS in a different storage buffer formulation.