

For Research Use Only

RPAIN Monoclonal antibody

Catalog Number: 68546-1-Ig



Basic Information

Catalog Number: 68546-1-Ig	GenBank Accession Number: BC004451	Purification Method: Protein G purification
Size: 1000 µg/ml	GeneID (NCBI): 84268	CloneNo.: 2D8E3
Source: Mouse	UNIPROT ID: Q86UA6	Recommended Dilutions: WB 1:2000-1:10000
Isotype: IgG1	Full Name: RPA interacting protein	
Immunogen Catalog Number: AG7844	Calculated MW: 25 kDa	
	Observed MW: 25 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : A549 cells, Jurkat cells, A2780 cells, LNCap cells, K-562 cells
Species Specificity: Human	

Background Information

Replication protein A (RPA) is a eukaryotic single-stranded DNA binding protein that is essential for DNA replication, repair, and recombination, and human RPA interacting protein α (hRIP α) is the nuclear transporter of RPA. RPAIN was originally identified as a nuclear transporter of RPA in *Xenopus* (PMID: 23010595). The human RIP α gene encodes several splice isoforms, of which hRIP α and hRIP β are the major translation products in vivo. hRIP α transports RPA into the nucleus by shuttling between the cytoplasm and nucleus while hRIP β is localized in promyelocytic leukemia (PML) nuclear bodies.

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68546-1-Ig (RPA1N antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 68546-1-Ig (RPA1N antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.