## For Research Use Only

## **HELIOS Monoclonal antibody**

Catalog Number: 68558-1-Ig



**Purification Method:** 

**Basic Information** 

Catalog Number: GenBank Accession Number:

68558-1-Ig BC028936 Protein A purification Size: GeneID (NCBI): CloneNo.: 1000 μ g/ml 22807 1F1F10

Source: UNIPROT ID: Recommended Dilutions:

 Mouse
 Q9UKS7
 WB 1:2000-1:10000

 Isotype:
 Full Name:
 IP 0.5-4.0 ug for 1.0-3.0 mg of total

IgG2b IKAROS family zinc finger 2 (Helios) protein lysate

Immunogen Catalog Number:Calculated MW:AG4357500 aa, 57 kDa

Observed MW: 65 kDa

**Applications** 

Tested Applications: Positive Controls:

IP, WB, ELISA WB: Jurkat cells, MOLT-4 cells

Species Specificity: IP : Jurkat cells,

**Background Information** 

IKZF2 (Helios), a member of the Ikaros transcription factor family, is expressed in a majority of Treg cells and some T cell and lymphocyte subsets. The expression of IKZF2 is associated with T cell activation and cellular division and is suggested a suitable marker to distinguish natural and induced Treg (PMID:21918685). IKZF2 plays a crucial role in maintaining function and stability of Treg cells, and IKZF2 deficiency reduces tumor growth in mice (PMID: 36863346).

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



Jurkat cells were subjected to SDS PAGE followed by western blot with 68558-1-1g (HELIOS antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



IP result of anti-HELIOS (IP:68558-1-Ig, 4ug; Detection:68558-1-Ig 1:3000) with Jurkat cells lysate 1560 ug.