For Research Use Only

BCOR Monoclonal antibody, PBS Only proteintech

Catalog Number: 68611-1-PBS



Purification Method:

Protein G purification

CloneNo.:

1F12H6

Basic Information

Catalog Number:

Size: 1 mg/ml

68611-1-PBS BC009675

GeneID (NCBI): 54880

GenBank Accession Number:

Source: UNIPROT ID: Mouse Q6W2J9
Isotype: Full Name:

IgG1 BCL6 co-repressor
Immunogen Catalog Number: Calculated MW:

AG33785 1703 aa, 186 kDa Observed MW:

Observed MV 186-200 kDa

Applications

Tested Applications: WB,Indirect ELISA Species Specificity: Human, mouse

Background Information

BCOR, a corepressor of BCL6, binds to transcriptional factors and specifically inhibits gene expression. BCOR plays a key role in the early embryonic development, mesenchymal stem cell function and hematopiesis. BCOR can associate with HDACs, the polycomb group protein, polycomb group Ring finger protein1/nervous system polycomb1 (PCGF1/NSPC1), histone demethylase, F-box- and leucine-rich repeat protein10(FBXL10). Via repression of TFAP2A acts as a negative regulator of osteo-dentiogenic capacity in adult stem.

Storage

Storage: Store at -80°C.


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer: PBS Only

Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 68611-1-1g (BCOR antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 68611-1-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 68611-1-1g (BCOR antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 68611-1-PBS in a different storage buffer formulation.



RAW 264.7 cells were subjected to SDS PAGE followed by western blot with 68611-1-1g (BCOR antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 68611-1-PBS in a different storage buffer formulation.



HEK-293 cells were subjected to SDS PAGE followed by western blot with 68611-1-1g (BCOR antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 68611-1-PBS in a different storage buffer formulation.