For Research Use Only

PRKCB Monoclonal antibody

Catalog Number:68620-1-lg 2 Publications



Basic Information

Applications

Catalog Number: 68620-1-lg

Size:
500 µg/ml
Source:
Mouse
Isotype:
IgG1

Immunogen Catalog Number:

AG33693

Tested Applications:

Cited Applications:

WB, IHC

WB, ELISA

Species Specificity: Human, pig, rabbit, rat, mouse

Cited Species:

rat

GenBank Accession Number:

BC036472 GeneID (NCBI): 5579 UNIPROT ID: P05771

Full Name: protein kinase C, beta

Calculated MW: 673 aa, 77 kDa Observed MW:

77 kDa

Positive Controls:

WB: human brain tissue, pig brain tissue, rabbit brain tissue, rat brain tissue, mouse brain tissue

Purification Method:

CloneNo.:

4B7A1

Protein G purification

Recommended Dilutions:

WB 1:2000-1:10000

Background Information

Protein kinase C (PKC) is a family of serine- and threonine-specific protein kinases that can be activated by calcium and second messenger diacylglycerol. Protein kinase Cs (PKCs) are signaling molecules that play key roles in many cellular processes, including secretion, gene expression, proliferation, and muscle contraction. The PKC family is broadly divided into three subgroups: classical, novel, and atypical PKCs. These subgroups differ in both protein sequences and mechanistic requirements for catalytic activity. PKC beta is one of the PKC family members. PKC beta has been reported to be involved in many different cellular functions, such as B cell activation, apoptosis induction, endothelial cell proliferation, and intestinal sugar absorption. Studies in mice also suggest that this kinase may also regulate neuronal functions and correlate fear-induced conflict behavior after stress. This antibody can recognize PKC beta, PKC alpha, PKC gamma, PKC epsilon and PKC delta.

Notable Publications

Author	Pubmed ID	Journal	Application
Kai Zhang	38718892	J Ethnopharmacol	WB,IHC
Xiaoling Li	38061206	Redox Biol	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68620-1-1g (PRKCB antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Alpha Tubulin Monoclonal antibody (HRP-66031) as loading control.