## For Research Use Only

## PARK7/DJ-1 Monoclonal antibody

Catalog Number: 68915-1-Ig



**Basic Information** 

Catalog Number:

GenBank Accession Number: BC008188

Purification Method: Protein G purification

68915-1-lg Size:

GeneID (NCBI): 11315 CloneNo.:

500 ug/ml Source: Mouse

UNIPROT ID: Q99497 4E10B2
Recommended Dilutions:

Isotype: IgG1 Full Name: Parkinson disease (autosomal WB 1:2000-1:10000

Immunogen Catalog Number: AG28526

recessive, early onset) 7
Calculated MW:

189 aa, 20 kDa

Observed MW: 20-25 kDa

**Applications** 

**Tested Applications:** 

WB, ELISA

Species Specificity: human, pig, rabbit **Positive Controls:** 

WB: HeLa cells, pig brain tissue, HEK-293 cells, Jurkat

cells, rabbit brain tissue

## **Background Information**

PARK7, also named as DJ1, belongs to the peptidase C56 family. It protects cells against oxidative stress and cell death. PARK7 plays a role in regulating expression or stability of the mitochondrial uncoupling proteins SLC25A14 and SLC25A27 in dopaminergic neurons of the substantia nigra pars compacta and attenuates the oxidative stress induced by calcium entry into the neurons via L-type channels during pacemaking. It eliminates hydrogen peroxide and protects cells against hydrogen peroxide-induced cell death. PARK7 has cell-growth promoting activity and transforming activity. It may function as a redox-sensitive chaperone. It's precursor undergoes a cleavage of a C-terminal peptide and subsequent activation of protease activity in response to oxidative stress. The amino acid replace at 166 (L — P) reduces PARK7 protein stability and leads to increased degradation. The predicted MW of this protein is 20 kDa, An additional 25 kDa band can be observed due to modification (PMID: 31767755).

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68915-1-1g (PARK7/DJ-1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 68915-1-1g (PARK7/DJ-1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.