For Research Use Only

NeutraControl EPO Mouse McAb

Catalog Number:69519-1-Ig





Basic Information	Catalog Number: 69519-1-Ig Concentration: Source: Mouse Isotype: IgG1 Immunogen Catalog Number: HZ-1168	GenBank Accession Number: GeneID (NCBI): 2056 Full Name: erythropoietin	Purification Method: Protein G purification CloneNo.: 1D3B1
Applications	Tested Applications: ELISA, Non-Neutralization Species Specificity: human		trols: ization : TF-1 cells, mbinant protein,
Background Information	Erythropoietin (Epo) is a member of the EPO/TPO family and encodes a secreted, glycosylated cytokine composed of four alpha helical bundles. The protein is found in the plasma and regulates red cell production by promoting erythroid differentiation and initiating hemoglobin synthesis. Its effect is realized by binding erythropoietin receptor (EpoR) expressed on erythroid progenitor cells. EpoR, is a glycoprotein expressed on megakaryocytes, erythroid progenitors and endothelial cells. Epo also has neuroprotective activity against a variety of potential brain injuries and antiapoptotic functions in several tissue types. This antibody is a neutralizing control antibody for EPO, the immunogen is the same as Neutrakine 69019-1-lg but could not neutralize human EPO.		
Storage	Storage: Lyophilized antibodies are stable for 1 year from the date of receipt if stored between (-20°C) and (-80°C). Upon reconstitution we recommend that the solution can be stored at(4°C) for short term or at(-20°C) to (-80°C) for long term. Repeated freeze thaw cycles should be avoided with reconstituted products. Storage Buffer: Sterile PBS, pH7.4 Aliquoting is unnecessary for -20°C storage		

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Recombinant human EPO (Cat.NO. HZ-1168) stimulates TF-1 cells (human erythroleukemic cell line) in a dose-dependent nammer (blue curve, refer to bottom X-left Y). The activity of human EPO (1 ng/mL HZ-1168) was neutralized by mouse anti-human EPO monoclonal antibody 69019-1-Ig at serial dose (red curve, refer to top X-right Y). The ND50 is typically 2-5 µ g/mL. The NeutraControl mouse anti-human EPO monoclonal antibody 69519-1-Ig could

Indirect ELISA was carried out by coating recombinant Human EPO (Cat.NO. HZ-1168) at 70 ng/well followed by blocking and adding serial diluted EPO antibody 69019-1-Ig and 69519-1-Ig respectively. Signal was developed with TMB and stopped by H2SO4. Signal strength was measured by absorbance at 450 nm.