

For Research Use Only

# Phospho-HSP27 (Ser78) Recombinant antibody

Catalog Number: 80046-3-RR



## Basic Information

**Catalog Number:**

80046-3-RR

**Size:**

1000 µg/ml

**Source:**

Rabbit

**Isotype:**

IgG

**GenBank Accession Number:**

BC012768

**GeneID (NCBI):**

3315

**UNIPROT ID:**

P04792

**Full Name:**

heat shock 27kDa protein 1

**Calculated MW:**

19-23 kDa

**Observed MW:**

25-30 kDa

**Purification Method:**

Protein A purification

**CloneNo.:**

241528D11

**Recommended Dilutions:**

WB 1:5000-1:50000

## Applications

**Tested Applications:**

WB, ELISA

**Species Specificity:**

human

**Positive Controls:**

WB : Calyculin A treated PC-3 cells,

## Background Information

HSPB1, also named as HSP27, belongs to the mammalian small HSP family widely expressed in cells including the cytoplasm and nucleus. It has been reported that HSPB1 plays a key role in stress resistance, actin organization and numerous biological processes. HSPB1 can be phosphorylated in different stimulation such as cytokines, growth factors and peptide hormones on the site of Ser15, Ser78 and Ser82 by MAPKAP2 kinase. The phosphorylation of HSPB1 leads to a decrease in the size of HSPB1 oligomers. HSPB1 is associated with amyotrophic lateral sclerosis (ALS) (PMID: 8542580, 11960785, 10383393, 32231288)

## Storage

**Storage:**

Store at -20°C. Stable for one year after shipment.

**Storage Buffer:**

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Non treated and Calyculin A treated PC-3 cells were subjected to SDS PAGE followed by western blot with 80046-3-RR (Phospho-HSP27 (Ser78) antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with Alpha Tubulin (66031-1-Ig) antibody as a loading control.