

For Research Use Only

Phospho-PRKD1 (Ser916) Recombinant antibody, PBS Only

Catalog Number:80080-2-PBS



Basic Information

Catalog Number: 80080-2-PBS	GenBank Accession Number: NM_001330069	Purification Method: Protein A purification
Size: 1 mg/ml	GeneID (NCBI): 5587	CloneNo.: 241786A11
Source: Rabbit	UNIPROT ID: Q15139	
Isotype: IgG	Full Name: protein kinase D1	
	Calculated MW: 102 kDa	
	Observed MW: 115 kDa	

Applications

Tested Applications:
WB, Indirect ELISA

Species Specificity:
human, mouse

Background Information

Protein kinase D1 (PRKD1), also named as PKD1 and PKC μ , is comprised of two cysteine-rich domains and a pleckstrin homology (PH) domain. PKD1 is involved in cellular processes including protein secretion, proliferation, cytoskeletal reorganization, Golgi function, immune function and apoptosis. It is widely expressed in thyroid, brain, heart, lung and other tissues. PKCs have been shown to regulate PKD1 activation. It has been reported that ser 916 is a PKD1 autophosphorylation site. PKD1 can be activated by growth factors, oxidative stress, thrombin, bioactive lipids, cross-linking of B- and T-cell receptors and some G-protein coupled receptors (GPCR). PKD1 is located mainly in the cytoplasm in unstimulated cells, while PKD1 migrates to the membrane in activated cells. (PMID: 17306383, 24806360, 30101477, 21696630)

Storage

Storage:
Store at -80°C.
The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:
PBS Only

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Non-treated NIH/3T3 cells, phosphatase inhibitor treated NIH/3T3 cells and λ phosphatase treated NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 80080-2-RR (Phospho-PRKD1 (Ser916) antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with Vinculin (66305-1-Ig) antibody as a loading control. This data was developed using the same antibody clone with 80080-2-

