For Research Use Only

## Phospho-JUN (Ser73) Recombinant antibody, PBS Only



Catalog Number:80086-1-PBS

**Basic Information** 

Catalog Number: 80086-1-PBS

Size: 1 mg/ml Source: Rabbit Isotype: GenBank Accession Number:

BC068522 GeneID (NCBI): 3725 UNIPROT ID: P05412 Full Name: jun oncogene

Calculated MW: 331 aa, 36 kDa Observed MW: 42-45 kDa Purification Method: Protein A purification

CloneNo.: 4A18

**Applications** 

Tested Applications: WB,Indirect ELISA Species Specificity: Human, mouse


## **Background Information**

JUN, the most extensively studied protein of the activator protein-1 (AP-1) complex, is involved in numerous cell activities, such as proliferation, apoptosis, survival, tumorigenesis and tissue morphogenesis (PMID: 22180088). JUN is a transcription factor that recognizes and binds to the enhancer heptamer motif 5'-TGA[CG]TCA-3'. It promotes activity of NR5A1 when phosphorylated by HIPK3 leading to increased steroidogenic gene expression upon cAMP signaling pathway stimulation. JUN is a basic leucine zipper (bZIP) transcription factor that acts as homo- or heterodimer, binding to DNA and regulating gene transcription (PMID: 9732876). In additon, extracellular signals can induce post-translational modifications of JUN, resulting in altered transcriptional activity and target gene expression (PMID:8464713). More over, it has uncovered multiple layers of a complex regulatory scheme in which JUN is able to crosstalk, amplify and integrate different signals for tissue development and disease. Jun is predominantly nuclear, ubiquitinated Jun colocalizes with lysosomal proteins (PMID: 15469925). This antibody is raised against synthetic phosphopeptide corresponding to residues surrounding Ser73 of human JUN, which can detect the bands around 42-45 kDa.

Storage

Storage: Store at -80°C. Storage Buffer: PBS Only

## **Selected Validation Data**



UV treated and non-treated NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 80086-1-RR (Phospho-JUN (Ser73) antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with Beta Actin antibody (66009-1-1g) as loading control. This data was developed using the same antibody clone with 80086-1-PBS in a different storage buffer formulation.